

Annexin A1(C5F8),MMab

Mouse Monoclonal

【Catalog Number】

REF R-221-2

【Package Size】

Ready to use: ☐ 1mL ☐ 2mL ☐ 3mL ☐ 5mL ☐ 6mL

Concentrated: ☐ 0.1mL ☐ 0.2 mL ☐ 0.5mL ☐ 1.0mL

【Intended Use】

For in vitro diagnostic use. This antibody is intended for immunohistochemical analysis on formalin-fixed, paraffin-embedded (FFPE) tissues. Results should be interpreted by a qualified pathologist using appropriate controls and relevant clinical information.

This antibody is for research use only (RUO).

【Specimen Collection and Preparation for Analysis】

Formalin-fixed, paraffin-embedded tissues.

Each section should be cut to the appropriate thickness (2-5 μ m) for the primary antibody being used and placed on a positively charged glass microscope slide.

【Storage and Handling】

Store at 2-8°C. Do not freeze.

Do not use product beyond the expiration date for the storage method.

【Reagents Provided】

Clone: C5F8

Buffer: 10mM pH 7.4 Phosphate Puffer Saline (PBS).

Stabilizer: 0.05% bovine serum (BSA).

Preservative: 0.05% sodium azide (NaN_3).

Ready-to-use antibody concentration: 2-5 μ g/mL.

Concentrated antibody concentration: 50-200 μ g/mL.

【Staining Procedure】

1. Deparaffinized slides in 3 changes of xylene (or Dewax solution), 10 minutes each. and hydrate through a graded series alcohols .

2. Wash the section in 90%, 80% and 70% ethyl alcohol for 10 minutes each.

3. Rinse in distilled water, 2 x 5 minutes.

4. Block the endogenous peroxidase by incubating the tissue in 3% hydrogen peroxide (H_2O_2) for 10 minutes.

5. Wash in distilled water, 2 x 5 minutes.

6. Antigen retrieval: Place slides in a pressure cooker filled with Epitope Retrieval Solution (Citrate, pH 6.0) buffer.

7. Wash in PBS 2 x 5 minutes.

8. Concentrated Antibody Dilution

Suggested Dilution: 1:50-1:100

The optimal dilution for a specific application under a given set of experimental conditions should be determined by the investigator.

9. Add 100 μ L primary antibody, Incubate for 30 minutes . Wash in PBS 2 x 5 minutes.

10. Add 100 μ L secondary antibody (the protocol depends on the supplier), and proceed to standard immunohistochemistry protocol. Wash in PBS 2 x 5 minutes.

11. Add 100 μ L DAB solution (the protocol depends on the supplier), Incubate for 2-10 minutes. Wash in PBS 2 x 5 minutes.

12. Counterstain with hematoxylin. Rinse with deionized water.

Application

For IVD purposes should be test and confirmed by lab (RUO)

【Contact Information】



Euromab S.r.l.

Add: Via Tortona, 21

20144 Milano (MI)

Italy

Email: info@euromab.com

Web: www.euromab.com