

## CD23 (C3B11), MMab

Mouse Monoclonal

### 【Catalog Number】

REF R-483-2

### 【Package Size】

Ready to use:  1mL  2mL  3mL  5mL  6mL

Concentrated:  0.1mL  0.2 mL  0.5mL  1.0mL

### 【Intended Use】

This mouse monoclonal antibody (Clone C3B11) is designed for immunohistochemical detection of CD23 antigen on formalin-fixed, paraffin-embedded human tissues. It is intended for in vitro diagnostic use and should be interpreted by a qualified pathologist in conjunction with clinical and morphological data.

***This antibody is for research use only (RUO).***

### 【Specimen Collection and Preparation for Analysis】

Formalin-fixed, paraffin-embedded tissues.

Each section should be cut to the appropriate thickness (2-5 µm) for the primary antibody being used and placed on a positively charged glass microscope slide.

### 【Storage and Handling】

Store at 2-8°C. Do not freeze.

Do not use product beyond the expiration date for the storage method.

### 【Reagents Provided】

#### **Clone: C3B11**

Buffer: 10mM pH 7.4 Phosphate Puffer Saline (PBS).

Stabilizer: 0.05% bovine serum (BSA).

Preservative: 0.05% sodium azide (NaN<sub>3</sub>).

Ready-to-use antibody concentration: 2-5µg/mL.

Concentrated antibody concentration: 50-200µg/mL.

### 【Staining Procedure】

1. Deparaffinized slides in 3 changes of xylene (or Dewax solution), 10 minutes each. and hydrate through a graded series alcohols .

2. Wash the section in 90%, 80% and 70% ethyl alcohol for 10 minutes each.

3. Rinse in distilled water, 2 x 5 minutes.

4. Block the endogenous peroxidase by incubating the tissue in 3% hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) for 10 minutes.

5. Wash in distilled water, 2 x 5 minutes.

6. Antigen retrieval: Place slides in a pressure cooker filled with Epitope Retrieval Solution (Citrate, pH 6.0) buffer.

7. Wash in PBS 2 x 5 minutes.

#### 8. Concentrated Antibody Dilution

Suggested Dilution: 1:50-1:100

The optimal dilution for a specific application under a given set of experimental conditions should be determined by the investigator.

9. Add 100µL primary antibody, Incubate for 30 minutes . Wash in PBS 2 x 5 minutes.

10. Add 100µL secondary antibody (the protocol depends on the supplier), and proceed to standard immunohistochemistry protocol. Wash in PBS 2 x 5 minutes.

11. Add 100µL DAB solution (the protocol depends on the supplier), Incubate for 2-10 minutes. Wash in PBS 2 x 5 minutes.

12. Counterstain with hematoxylin. Rinse with deionized water.

### **Application**

*For IVD purposes should be test and confirmed by lab (RUO)*

### 【Contact Information】



#### **Euromab S.r.l.**

Add: Via Tortona, 21

20144 Milano (MI)

Italy

Email: [info@euromab.com](mailto:info@euromab.com)

Web: [www.euromab.com](http://www.euromab.com)